Silica-Sand Deposits in Arizona

Jon E. Spencer and Nyal Niemuth
Arizona Geological Survey

OPEN-FILE REPORT OFR-15-04

August 2015

Arizona Geological Survey

www.azgs.az.gov | repository.azgs.az.gov

Arizona Geological Survey

M. Lee Allison, State Geologist and Director

Manuscript approved for publication in August 2015
Printed by the Arizona Geological Survey
All rights reserved

For an electronic copy of this publication: www.repository.azgs.az.gov
Printed copies are on sale at the Arizona Experience Store
416 W. Congress, Tucson, AZ 85701 (520.770.3500)

For information on the mission, objectives or geologic products of the
Arizona Geological Survey visit www.azgs.az.gov.

This publication was prepared by an agency of the State of Arizona. The State of Arizona, or any agency thereof, or any of their employees, makes no warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed in this report. Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the State of Arizona.

Recommended Citation: Spencer, J.E. and Niemuth, N.J., 2015, Silica-sand deposits in Arizona. Arizona Geological Survey Open File Report, OFR-15-04, 8 p.

Silica-sand deposits in Arizona

Jon E. Spencer
Arizona Geological Survey
416 W. Congress St., #100
Tucson, AZ 85701
jon.spencer@azgs.az.gov

Nyal J. Niemuth
Arizona Geological Survey, Phoenix Branch Office
3550 N Central Ave., 2nd floor
Phoenix, AZ 85012
nyal.niemuth@azgs.az.gov

Introduction

Nearly pure quartz sands have become increasingly important for oil and gas production. Although used for hydraulic fracturing ("fracking") since at least the 1950s, the advent of directional drilling and improvements in hydraulic fracturing techniques have led to widespread exploitation of quartz sands that have appropriate properties for use in hydraulic fracturing ("frac sands"). Such sands are present in the upper Miocene to Pliocene upper Bidahochi Formation of northeastern Arizona. Other sand deposits are present in the State, but are not as well suited for use in hydraulic fracturing. This report is a brief review of the distribution and character of sands in Arizona that may have properties appropriate for use in hydraulic fracturing.

Bidahochi Formation

The Bidahochi Formation in northeastern Arizona (Figures 1, 2) consists of three members: (1) a lower playa and lacustrine facies of middle Miocene age (Dallegge et al., 2001, 2003), (2) a middle member consisting of mafic lava, volcaniclastic sandstone, and tuff of upper Miocene age (Sutton, 1974; Hackman and Olson, 1977; Ulrich et al., 1984; Damon and Spencer, 2001), and (3) an upper member that consists of fluvial and eolian sand, silt, and clay of uppermost Miocene to Pliocene age (Repenning and Irwin, 1954; Repenning et al., 1958; Love, 1989). Although the Bidahochi Formation has been subdivided into members, the members are facies with complex depositional relationships (e.g., Kiersch and Keller, 1955; Dickinson, 2013). The lower and middle members are completely missing from southern and eastern areas where the upper member rests on Paleozoic and Mesozoic strata (Figure 2).

In the Sanders area the upper, dominantly fluvial member includes bentonite clay deposits (it is not entirely clear that assignment of Bidahochi Formation members is meaningful in this area). The bentonite clay deposits, consisting of calcium montmorillonite and known as the Cheto mine deposits, have been exploited for desiccants, gellants, and acid-activated bentonites since 1924 (Kiersch and Keller, 1955; Eyde and Eyde, 1987). The overlying silts and sands of the upper Bidahochi Formation include sand deposits that contain 93-99% SiO_2 (Table 1), some of which contain a large fraction of sand in the -20 +40 mesh size that is generally preferred for frac sands (Tables 2-5). Furthermore, the deposits can be mined with a front-end loader, unlike other frac-sand deposits in Texas and elsewhere that must be mined and processed by blasting and crushing (Eyde and Eyde, 1987).

Figure 1. Geologic map of Arizona (from Richard et al., 2000) showing locations of sand-dune deposits in western Arizona and location of Figure 2.

The Preferred Sands plant and quarry (Figure 3) were active in 2015, with employment of 53 workers according to the U.S. Department of Labor (http://www.msha.gov/drs/drshome.htm). Employment reached 95 workers in 2012 (actually "sum of average employment" based on hours worked as reported to the USDOL). The quarry operations are located largely at the site of the former Cheto bentonite quarries. Arizona Silica Sand Company was engaged in sand production at Houck through 2011 according to the Arizona Mine Inspector's annual report. However, the Houck silica sand plant and quarry (Figure 3) appeared from aerial imagery to be inactive, and a search of the U.S. Department of Labor web site yielded no information using the search terms "Arizona Silica", "Houck Silica" and "Houck".



Modified from figure 2 *in* Dickinson, 2013, Geosphere, v. 9, p. 1–20

Figure 2. Distribution of the Bidahochi Formation in northeastern Arizona and adjacent New Mexico (modified from Dickinson, 2013). Also shown is a simplified outline of the extent of the Navajo Indian Reservation. In some areas mineral rights and surface rights are held by different parties.

Lower Colorado River Valley

Sand dunes are exposed over large areas in the Yuma Desert southeast of Yuma and at Cactus Plain southeast of Parker (Figure 1). In both areas sands deposited in the Colorado River Valley by the Colorado River have been displaced by winds and redeposited in sand dunes. Chemical analysis of dune sands from Cactus Plain near Parker and the Algodones Dunes in the Yuma desert near Yuma indicate silica content primarily in the 80-90% range, with none exceeding 90% (Zimbelman and Williams, 2002). Analyses of dune sands in the eastern Mojave Desert, west of the Colorado River, indicate silica content of consistently <80%, apparently because these sand deposits are upwind rather than downwind from quartz-rich sands delivered by the Colorado River (Zimbelman and Williams, 2002). Although Cactus Plain and the Algodones Dunes are both near railroad tracks and could be easily quarried, their physical properties are not appropriate for hydraulic fracturing sands because they contain too much feldspar and perhaps because of other factors such as small grain size.

Conclusion

The unusual purity of sand deposits in the upper Bidahochi Formation, containing little but quartz, and the adequate roundness and grain size, have made these deposits attractive for use in hydraulic fracturing. Other deposits in Arizona that could be economically quarried and processed have not been identified. Given the enormous aerial extent of the upper Bidahochi Formation, it seems likely that this area will continue to be exploited for frac sands as long as hydraulic fracturing remains a viable tool for oil and gas extraction.

Table 1. Geochemical analysis of sands (from unpublished AZGS file data)

	Arizona Silica Sand Co., 1992	Arizona Silica in Houck (probably Arizona Silica Sand Company) 1989 (12/20 mesh)	Arizona Silica in Houck (probably Arizona Silica Sand Company) 1989 (20/40 mesh)	Arizona Silica in Houck (probably Arizona Silica Sand Company) 1989 (-40 mesh)	Navajo sand deposit north of Chambers (1956)
SiO_2	93	98.6	98.5	96.0	97.91
Al_2O_3	1.4	0.5	0.56	2.74	0.94
Fe_2O_3	1.9	0.108	0.12	0.32	0.14
CaO	0.09	0.12	0.14	0.22	0.12
Na_2O	0.24				0.23
K_2O	0.7				0.09
MgO	0.08	<0.1%	<0.01	<0.01	0.08
TiO_2	0.02	0.030	0.030	0.026	
LOI (volitiles lost upon heating to 1000°C)*	0.2	0.61	0.61	0.64	0.94
Total	97.63	99.968	99.960	99.946	100.45

*LOI represents % acid-soluble material

Table 2. Sand seive size (% retained) (from unpublished AZGS file data)

Seive size	Arizona Silica Sand Company, 1993	Arizona Silica Sand Company, 1993 (plaster sand)
+30	0	1.2
-30 +40	19.8	6
-40 +50	46.7	24.2
-50 +60	13.6	12.3
-60 +70	9.5	14.1
-70 +100	9.4	25.7
pan	1	16.5
total	100	100

Table 3. Sand seive size (% retained) (from unpublished AZGS file data)

Seive size	Balcomb lease in Burntwater Wash (76% of sands -10 +60)
-10 +20	15.7
-20 +40	39.3
-40 +60	21.4

Table 4. Sand seive size (% retained), sand deposit north of Chambers, location uncertain (from unpublished 1956 AZGS file data)

Seive size	Area 1	Area 2	Area 3	Area 4	Area 5
+10	0.11	0.11	0.04	1.05	1.27
-10 +20	35.31	37.2	29.53	38.17	27.09
-20 +40	32	39.34	34.34	21.28	28.39
-40 +60	16.5	12.2	21.68	13.9	19.58
-60	16.08	11.15	14.41	25.6	23.67

Table 5. Sand seive size (% retained)(from Kiersch, 1955)

Seive size	Bidahochi Fm., 14 miles N of Chambers	Bidahochi Fm., 1/2 mile SW of Klagetoh (~20 miles N of Chambers)
-12 +16	34.1	0.1
-16 +20	61.1	16.7
-20 +32	2.8	40.1
-32 +40	0.4	30.5
-40 +60	0.3	6.7
-60	0.8	5.3
total	99.5	99.4



Figure 3. Location map for quarries and facilities associated with silica sand deposits in the Sanders – Houck area.

References Cited

Dallegge, T.A., Ort, M.H., and McIntosh, W.C., 2003, Mio-Pliocene chronostratigraphy, basin morphology and paleodrainage relations derived from the Bidahochi Formation, Hopi and Navajo Nations, northeastern Arizona**:** The Mountain Geologist, v. 40, n. 3, p. 55-82.

Dallegge, T.A., Ort, M.H., McIntosh, W.C., and Perkins, M.E., 2001, Age and depositional basin morphology of the Bidahochi Formation and implications for the ancestral Upper Colorado River, *in* Young, R.A., and Spamer, E.E., eds., The Colorado River: Origin and evolution: Grand Canyon, Arizona, Grand Canyon Association Monograph 12, p. 47-51.

Damon, P.E., and Spencer, J.E., 2001, K-Ar geochronologic survey of the Hopi Buttes volcanic field, *in* Young, R.A., and Spamer, E.E., eds., The Colorado River: Origin and evolution: Grand Canyon, Arizona, Grand Canyon Association Monograph 12, p. 53-56.

Dickinson, W.R., 2013, Rejection of the lake spillover model for initial incision of the Grand Canyon, and discussion of alternatives: Geosphere, v. 9, n. 1, p. 1–20; doi:10.1130/GES00839.1.

Eyde, T.H., and Eyde, D.T., 1987, Bentonite and specialty sand deposits in the Bidahochi Formation, *in* Peirce, H.W., ed., 21st Forum on the geology of industrial minerals: Arizona Bureau of Geology and Mineral Technology (now Arizona Geological Survey), Special Paper 4, p. 123-127.

Hackman, R.J., and Olson, A.B., compilers, 1977, Geology, structure, and uranium deposits of the Gallup 1° x 2° Quadrangle, New Mexico and Arizona: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-981, scale 1:250,000.

Kiersch, G.A., 1955, Mineral resources, Navajo-Hopi Reservations, Arizona-Utah, Volume II, Nonmetallic minerals - Geology, evaluation, and uses, with sections by J.W. Anthony, R.L. DuBois, P.W. Howell, J.C. Haff, W.D.Keller, H.E. Krumlauf, and R.L. Wilson: Tucson, University of Arizona Press, 105 p.

Kiersch, G.A., and Keller, W.D., 1955, Bleaching clay deposits, Sanders-Defiance Plateau District, Navajo country, Arizona: Society of Economic Geologists, Economic Geology, v. 50, p. 469-494.

Love, D.W., 1989, Bidahochi Formation: An interpretive summary, *in* Anderson, O.J., Lucas, S.G., Love, D.W., and Cather, S.M., eds., Southeastern Colorado Plateau: New Mexico Geological Society 40th Annual Field Conference Guidebook, p. 273–280.

Repenning, C.A., and Irwin , J.H., 1954, Bidahochi formation of Arizona and New Mexico: American Association of Petroleum Geologists Bulletin, v. 38, p. 1821-1826.

Repenning, C.A., Lance, J.F., Irwin, J.H., 1958, Tertiary stratigraphy of the Navajo Country, *in* Anderson, R.Y., and Harshbarger, J.W., eds., Guidebook of the Black Mesa Basin, northeastern Arizona: New Mexico Geological Society, Ninth Field Conference Guidebook, p. 123-129.

Richard, S.M., Reynolds, S.J., Spencer, J.E., and Pearthree, P.A., 2000, Geologic map of Arizona: Arizona Geological Survey Map 35, scale 1:1,000,000.

Sutton, R.L., 1974, Field guide for Hopi Buttes and Navajo Buttes areas, Arizona, *in* Karlstrom, T.N.V., Swann, G.A., and Eastwood, R.L., eds., Geology of northern Arizona with notes on archaeology and paleoclimate, Part 2 - Area Studies and Field Guides: Geological Society of America, Rocky Mountain Section, 27th Annual Meeting, Flagstaff, Arizona, p. 699-727.

Ulrich, G.E., Billingsley, G.H., Hereford, R., Wolfe, E.W., Nealey, L.D., and Sutton, R.L., compilers, 1984, Geology, structure, and uranium deposits of the Flagstaff 1° x 2° Quadrangle, Arizona: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-1446, scale 1:250,000.

Zimbelman, J.R., and Williams, S.H., 2002, Geochemical indicators of separate sources for eolian sands in the eastern Mojave Desert, California, and western Arizona: Geological Society of America Bulletin, v. 114, n. 4, p. 490-496.